

GREY BRUCE HEALTH UNIT
MOSQUITO IDENTIFICATION AND VIRAL TESTING
SUMMARY REPORT
2018

SPOROMETRICS
219 DUFFERIN STREET SUITE 20C
TORONTO, ONTARIO M6K 1Y9

t. 416 516 1660
f. 416 516 1670
www.sporometrics.com



Sporometrics

TABLE OF CONTENTS

EXECUTIVE SUMMARY	II
1.0 INTRODUCTION.....	1
2.0 BACKGROUND.....	1
3.0 SCOPE OF WORK	2
3.1 Overall Tasks	2
3.2 Mosquito Species Identification	2
3.3 Viral Testing	3
4.0 SUMMARY OF 2018 RESULTS	4
4.1 Vector Abundance.....	4
4.2 Summary of Traps.....	5
4.3 Summary of Viral Testing Pools	6
4.4 Accumulated Degree Days.....	9
5.0 DISCUSSION.....	9
6.0 CLOSURE.....	10

References

EXECUTIVE SUMMARY

Sporometrics performed mosquito identification and laboratory analysis for the Grey Bruce Health Unit (GBHU) for the 2018 West Nile Virus (WNv) and Eastern Equine Encephalitis Virus (EEEv) mosquito surveillance program. Traps were received by Sporometrics from May 16th, 2018 (week 20) to September 26th, 2018 (week 39). In total 263 traps were sent, with approximately 12-17 traps sent every week for 20 weeks.

Mosquitoes were identified to a maximum of 150 per trap. Three pools (maximum 50 mosquitoes per pool) were tested for WNv and/or EEEv in accordance with the Ministry of Health and Long-Term Care (MOHLTC) directive. The insects were homogenized and processed through RNA extraction which was used as a template for qRT-PCR reactions. Only female mosquitoes of vector species were tested using the molecular assay.

The WNv testing priority in descending order was *Cx. pipiens / restuans*, *Cx. salinarius*, *Oc. japonicas*, *Cx. tarsalis*, *Ae. vexans vexans*, *Oc. triseriatus*, *An. punctipennis*, *Oc. trivittatus*, *An. walkeri* and *Oc. stimulans*. The EEEv testing priority in descending order was *Cs. melanura*, *Oc. canadensis*, *Cq. perturbans* and *Ae. vexans vexans*.

Throughout the season 88,894 mosquitoes were collected and 11,146 were identified. A total of 561 molecular tests were performed. Of these, 276 pools comprising 1,132 specimens were tested for WNv and 285 pools with 4,149 specimens were tested for EEEv. There was a positive pool found for WNv.

1.0 INTRODUCTION

Sporometrics Inc. (Sporometrics) is pleased to provide the Grey Bruce Health Unit (GBHU) with this Summary Report on the 2018 Mosquito Identification and Viral Testing program.

The 2018 Mosquito Identification and Viral Testing program included data management and reporting, accurate mosquito sorting, identification to the species level and viral testing of identified mosquito pools by qRT-PCR with generic and envelope TaqMan PCR assays. These viral testing protocols conformed to, and in some cases exceeded, the requirements of the National Steering Committee on West Nile virus Surveillance and the Ministry of Health and Long-Term Care's (MOHLTC) Gold Standard. Weekly results were submitted to the GBHU and the MOHLTC in the form of Microsoft Excel spreadsheets.

2.0 BACKGROUND

West Nile Virus (WNv; family *Flaviviridae*, genus *Flavivirus*) is an arthropod-borne virus or arbovirus that was originally isolated in the West Nile region of Uganda in 1937 (Smithburn et al. 1940). Since then, it has spread around the globe, sparing only Antarctica (Weaver et al. 2004, Kramer et al. 2008). It is still unknown how the virus came to North America, but WNv first appeared in New York City in 1999 (Lanciotti et al. 1999; Nash et al. 2001). Soon after, the virus spread to the rest of the continent (Weaver et al. 2004). WNv is maintained in a cycle involving mosquitoes and birds: virus-naïve mosquitoes feed on WNv-infected birds and become vectors for the virus, capable of continuing the cycle by biting other birds or acting as bridge vectors by biting incidental hosts such as humans and horses (Go et al. 2014).

Approximately 80% of all cases of human WNv infections are asymptomatic. The remaining 20% may develop fever, head and body aches, skin rash, generalized weakness, chills, joint pain, and/or painful eyes (Gray et al. 2014; Mostashari et al. 2001; Solomon 2004; Hubalek et al. 2001). The virus can cross the blood-brain barrier in less than 1% of cases, leading to meningitis, encephalitis, or flaccid paralysis (Kramer 2007).

WNv was first detected in Ontario in 2001 in birds, and in the following year, 395 confirmed and probable human cases were recorded (PHO 2012 report; PHO 2013 guide). The prevalence of human cases has fluctuated from 4 in 2009 to 269 in 2012 to 52 in 2013 (PHO 2012 report; NBPS 2014 report). Since the virus' appearance in Ontario, public health units have begun to trap mosquitoes in order to evaluate WNv risk based on the tendency of positive pools to occur in specific areas. In Ontario, more than 20 mosquito species have been found to be WNv-positive, despite the fact that *Culex pipiens* and *Cx. restuans* mosquitoes are the primary bridge vectors transmitting WNv to humans (PHO 2012 report; PHO 2008 preparedness plan). The weekly monitoring of these carriers is a high priority for public health officers, and rapid and accurate detection of carrier and vector identification is needed. Additional testing may be required if positive vectors attain unusual levels in any geographic areas within the region monitored.

Eastern Equine Encephalitis Virus (EEEV; family Togaviridae, genus Alphavirus) is also a mosquito-borne virus. It was first isolated in a horse in 1933 in New Jersey and Virginia (Hanson 1957; Scott et al. 1989). Like WNV, EEEV is maintained in a cycle between birds and mosquitoes, with bridge vectors transmitting the virus to incidental hosts like humans and horses (Go et al. 2014). Only a handful of human EEEV infections per year have been recorded since the 1960s (Go et al. 2014). While some cases are asymptomatic, EEEV is a health concern given that symptomatic patients usually develop severe encephalitis (Go et al. 2014). Not only is the fatality rate high (30 to 75%), EEEV survivors often develop neurological sequelae such as paralysis, brain dysfunction, and seizures (EEE report; Deresiewicz et al. 1997).

In Ontario, EEEV was first identified in mosquitoes in 2009 in a First Nations community within the Simcoe Muskoka District, but to date, no human infections have been identified (Parry Sound Report). The EEEV surveillance program began in 2010, focusing on *Cs. melanura*, *Oc. canadensis*, *Cq. perturbans* and *Ae. vexans vexans*, the vectors of EEEV (EEE 2014 report). The first Ontario public health unit to test EEEV-positive mosquitoes was the North Bay Parry Sound District in 2010 (EEE 2014 report; NBPS report). Between 2008 and 2013 inclusive, 14 equine cases and 16 EEEV-positive pools were reported in Ontario (EEE 2014 report). Monitoring for EEEV in bridge vectors has been mandated as a proactive approach to assessing the distribution of this emerging virus.

3.0 SCOPE OF WORK

The scope of work for this project was based on the GBHU Mosquito Identification and Viral Testing Program 2018 scope of work.

3.1 Overall Tasks

Trap receipt and sample reporting was conducted in accordance to Sporometrics standard operating procedures (SOPs) and ISO/IEC 17025:2005 practices. Data were collected on mosquito identification and abundance, and viral testing results for taxon pools as determined by MOHLTC guidelines. These data were compiled in a report data sheet, compatible with the MOHLTC and GBHU's software requirements, following Sporometrics SOPs for sample reporting. Sporometrics reported the data to the GBHU and MOHLTC each Monday following the receipt of the traps.

3.2 Mosquito Species Identification

Mosquitoes were pre-sorted by species and sex, and retained in pools of not more than 50 conspecific specimens at -80 °C prior to molecular analysis. Species identification was performed morphologically on the day the samples were received in the laboratory and according to accepted methods using standard references.

Sex differentiation

Males are readily differentiable from females of most mosquito species by the more elaborately ornamented, plumose morphology of their antennae. Compared to males, females also tend to have very short maxillary palpi relative to the proboscis. In males, the maxillary palpi tend to be longer than the proboscis.

Genus- and species-level identification

Identification of the adult mosquito is accomplished by examination of the insect by stereoscopic microscopy at magnifications up to 50x. Using stereomicroscopy alone, most of the important vectors of WNv / EEEv can be identified confidently to the level of species or species-complex by an experienced entomological technician using appropriate taxonomic references.

3.3 Viral Testing

Chain of custody

Sample receipt and reporting was conducted in accordance to Sporometrics SOPs and ISO/IEC 17025:2005 practices. Samples were either immediately processed for total RNA extraction and qRT-PCR analysis or stored at -80 °C until processing.

Mosquito extraction

Mosquito pools were selected and prioritized for WNv and/or EEEv testing based on MOHLTC guidelines. Sorted mosquitoes were pooled accordingly into polypropylene tubes and homogenized. The supernatant was used for viral RNA extraction.

WNv and EEEv testing

For the quantitative assessment of WNv the PCR protocol and cycling conditions followed accepted practices (Lanceotti et al. 2000). Samples were assessed by a qRT-PCR "TaqMan" assay that uses sets of primers and probes directed against the envelope and 3' non-coding region of the WNv genome that are more sensitive than traditional reverse transcriptase RT-PCR to identify WNv from positive mosquito pools (iScript One-Step RT-PCR Kit for Probes from Bio-Rad were used). TaqMan probes were labeled at the 5' end with the reporter dye FAM and at the 3' end with the quencher dye TAMRA (Applied Biosystems, Carlsbad, CA). Testing for EEEv followed the guidelines provided by the National Microbiology Laboratory (Public Health Agency of Canada, Winnipeg).

4.0 SUMMARY OF 2018 RESULTS

GBHU trapped mosquitoes every second week from week 24 to 34 (June 14th to August 23rd) during the season. The following sections summarize the results of the Mosquito Identification and Viral Testing program. A brief overview of the season's statistics is provided below:

Total number of traps	263
Total number of mosquitoes collected	88,894
Total number of mosquitoes identified	11,146
Number of pools viral tested for WNV	276
Number of WNV-positive pools	1
Number of pools viral tested for EEEV	285
Number of EEEV-positive pools	0

4.1 Vector Abundance

In 2018, the most abundantly captured species was *Cq. perturbans*, an EEEV vector. Of the 11,146 mosquitoes identified throughout the season, 3,443 were *Cq. perturbans*, representing 30.89% of identified species. The second most abundant species was *Oc. canadensis*, a WNV and EEEV vector, which comprised of 16.85% of the total amount of mosquitoes identified (11,146 mosquitoes). The population of all vector species is provided in Table 1 below.

Table 1 – Summary of identified vector species

Species	Quantity	Percentage (%)	Vector
<i>Cx. pipiens / restuans</i>	374	3.36%	WNV
<i>Cx. salinarius</i>	22	0.197%	WNV
<i>Oc. japonicus</i>	68	0.610%	WNV
<i>Cx. tarsalis</i>	0	0.00%	WNV
<i>Oc. triseriatus</i>	181	1.62%	WNV
<i>An. punctipennis</i>	252	2.26%	WNV
<i>Oc. trivittatus</i>	199	1.79%	WNV
<i>An. walkeri</i>	11	0.099%	WNV
<i>Oc. stimulans</i>	1,304	11.70%	WNV
<i>An. quadrimaculatus</i>	193	1.73%	WNV
<i>Ae. vexans vexans</i>	518	4.65%	WNV / EEEV
<i>Oc. canadensis</i>	1,878	16.85%	WNV / EEEV
<i>Cs. melanura</i>	0	0.00%	EEEV
<i>Cq. perturbans</i>	3,443	30.89%	EEEV
<i>Other</i>	291	2.61%	-
Total	11,146	100.00%	-

Table 1. Summary of vector species abundance and proportion relative to the taxa listed. Vector type for each species is indicated.

Cx. pipiens / restuans is a bridge vector and the primary vector for human cases of WNV in Ontario (Ontario mosquito guide). The following graph, Figure 1, illustrates the number of mosquitoes identified by week; the proportion of *Cx. pipiens / restuans*; and the total number of other WNV and EEEv vectors identified per week. The *Cx. pipiens / restuans* population was at its peak in week 34 and 35.

Figure 1 – Summary of identified vector species by week




Figure 1. Total identified vector populations from all traps by week. Stacked bar graph displays the total number of mosquitoes identified per week. Y-axis - number of mosquitoes, X-axis - epidemiological week number.

4.2 Summary of Traps

Traps SVCAB, HCBM and IPMK captured the most mosquitoes in the 2018 season, with 57,767, 11,111 and 8,701 mosquitoes, respectively. Trap HCBM had the most WNV vectors collected, trap SVCAB had the most WNv/EEEv vectors collected, and trap IPMK had the most EEEv vectors collected. Trap SVCAB also collected the highest quantity of *Cx. pipiens/restuans* (90 specimens) over the season. The following figure provides a summary of identified vector species by trap site.

Figure 2 – Summary of identified vector species by trap site




Figure 2. Total identified vector populations by trap. Stacked graph displays the total number of mosquitoes identified per trap site. Y-axis – number of mosquitoes; X-axis – trap site.

4.3 Summary of Viral Testing Pools

A total of 561 viral tests (consisting of 276 tests for WNV and 285 for EEEV) were performed throughout the season. One of the pools tested positive for WNV. The following tables provide a detailed overview of the pools tested.

Table 2a – Summary of WNv testing pools by species

WNv VT Summary by Species			
Species	Pools	Specimens	Positive Pools
<i>Cx. pipiens/restuans</i>	95	373	1
<i>Cx. salinarius</i>	5	13	0
<i>Oc. japonicus</i>	21	53	0
<i>Cx. tarsalis</i>	0	0	0
<i>Ae. vexans vexans</i>	51	232	0
<i>Oc. triseriatus</i>	15	53	0
<i>An. punctipennis</i>	32	55	0
<i>Oc. trivittatus</i>	13	65	0
<i>An. walkeri</i>	2	2	0
<i>Oc. stimulans</i>	28	229	0
<i>An. quadrimaculatus</i>	6	17	0
<i>Oc. canadensis</i>	8	40	0
Total	276	1132	1

Table 2a. WNv viral testing summary. Includes the number of pools tested, quantity of mosquitoes in the pools and the number of positives by species.

Table 2b – Summary of EEEv testing pools by species

EEEv VT Summary by Species			
Species	Pools	Specimens	Positive Pools
<i>Cs. melanura</i>	0	0	0
<i>Oc. canadensis</i>	88	1554	0
<i>Cq. perturbans</i>	108	2124	0
<i>Ae. vexans vexans</i>	89	471	0
Total	285	4149	0

Table 2b. EEEv viral testing summary. Includes the number of pools tested, quantity of mosquitoes in the pools and the number of positives by species.

A summary of the positive viral testing pool is provided in Table 3 below.

Table 3 – Summary of positive viral testing pool

Site Code	Week	Date Collected	Date Tested	Species	Number of mosquitoes tested	Number of mosquitoes collected	GPR1	EPR1	Final
PKMM	35	2018/08/29	2018/09/04	Cx. pipiens/restuans	2	2	33.38	36.73	Positive

Table 3. Summary of positive viral testing pools. Includes site code, species, generic primer region 1 (GPR1), and envelope primer region 1 (EPR1).

4.4 Accumulated Degree Days

A degree day is defined as a unit of measurement for temperature (PHO 2013 guide). It is the amount of heat an organism requires to develop through specific stages of their life cycle. A degree day is one day (24 hours) with a temperature below or above a fixed reference temperature. In vector surveillance, degree days are used to track when insects will proliferate. The MOHLTC uses 18.3°C for *Cx. pipiens / restuans*. Accumulated degree days (ADD) are the continuous addition of consecutive degree days from a set starting point.

In Ontario, WNV positive pools can occur as early as 30 ADD, and the number of positive pools can reach double-digit numbers once the ADD exceeds 140 (PHO 2013 guide). The first human cases are detected between 100 and 125 ADD (PHO 2013 guide). As illustrated in Figure 3 below, 30 ADD was reached at the end of week 26, 100 ADD was reached in week 30 and by the end of the trapping season, the ADD had reached 216.65.




Figure 3. Mean daily temperature and accumulated degree days for GBHU. Y-axis – mean temperature ($^{\circ}\text{C}$), X-axis – epidemiology week. Red line indicates daily mean temperature. Dashed line represents 18.3°C threshold. Orange shaded area represents accumulated degree days.

5.0 DISCUSSION

This year, the total number of mosquitoes identified was 11,146. This is an increase from the 2017 season. This year, the most abundantly trapped species was *Cq. perturbans* which is an EEEv vector. *Cq. perturbans* was identified in 30.89% of all identified species. In 2017, the most abundantly trapped species was *Ae. vexans vexans*. In 2017, *Ae. vexans vexans* was identified in 16.6% of all identified species. Vector abundance for *Cx. pipiens/restuans* has decreased from 2017. In 2018, only 3.36% of *Cx. pipiens/restuans* specimens were identified out of all identified species. Out of all mosquito species identified during the 2017 season, there were 9.32% *Cx. pipiens/restuans*.

One of the 276 pools tested for WNV was positive. After week 26, the ADD accumulated past 30, which suggests that positive WNV pools could be expected. Human cases may also be expected towards the middle of the season as the ADD exceeded 100 in week 30.

The data indicates that no change to the surveillance program is currently warranted.

6.0 CLOSURE

We trust that this summary report is in accordance with your requirements. Should you have any questions or require clarification on any element of this report, please feel free to contact the undersigned at any time.

Yours truly,

Sporometrics



James Scott, PhD, ARMCCM
CEO



Kristine White, B.A.
Project Manager

References

1. Armstrong PM, Andreadis TG. Eastern Equine Encephalitis virus in mosquitoes and their role as bridge vectors. *Emerg Infect Dis.* 2010;16(12):1869-1874.
2. Deresiewicz RL, Thaler SJ, Hsu L, et al. Clinical and neuroradiographic manifestations of eastern equine encephalitis. *N Eng J Med.* 1997;336(26):1867-1874.
3. Go YY, Balasuriya UBR, Lee C. Zoonotic encephalitides caused by arboviruses: transmission and epidemiology of alphaviruses and flaviviruses. *Clin Exp Vaccine Res.* 2014;3:58-77.

4. Gray TJ, Webb CE. A review of the epidemiological and clinical aspects of West Nile virus. *Int J Gen Med.* 2014;11(7):193-203.
5. Hanson RP. An epizootic of equine encephalomyelitis that occurred in Massachusetts in 1831. *Am J Trop Med Hyg.* 1957;6:858-862.
6. Hubalek Z. Comparative symptomatology of West Nile fever. *Lancet.* 2001;358(9278):254-255.
7. Kramer LD, Li J, Shi PY. West Nile virus. *Lancet Neurol.* 2007;6(2):171-181.
8. Kramer LD, Styler LM, Ebel GD. A global perspective on the epidemiology of West Nile virus. *Annu Rev Entomol.* 2008;53:61-81.
9. Lanciotti RS, Roehrig JT, Deubel V, et al. Origin of the West Nile virus responsible for an outbreak of encephalitis in the northeastern United States. *Science.* 1999;286(5448):2333-2337.
10. Lanciotti RS, Kerst AJ, Nasci RS, et al. Rapid detection of West Nile virus from human clinical specimens, field-collected mosquitoes, and avian samples by a TaqMan reverse transcriptase-PCR assay. *J Clin Microbiol.* 2000;38(11):4066-4071.
11. Mostashari F, Bunning ML, Kitsutani PT, et al. Epidemic West Nile encephalitis, New York, 1999: results of a household-based seroepidemiological survey. *Lancet.* 2001;358(9278):261-264.
12. Nash D, Mostashari F, Fine A, et al. The outbreak of West Nile virus infection in the New York City area in 1999. *N Engl J Med.* 2001;344:1807-1814.
13. North Bay Parry Sound District Health Unit (2014). 2014 Vector Borne Diseases Surveillance and Protection Plan. North Bay, ON: North Bay Parry Sound District Health Unit.
14. Ontario Agency for Health Protection and Promotion (Public Health Ontario). Eastern equine encephalitis: history and enhanced surveillance in Ontario. Toronto, ON: Queen's Printer for Ontario; 2014.
15. Ontario Agency for Health Protection and Promotion (Public Health Ontario). Guide for public health units: Considerations for adult mosquito control. Toronto, ON: Queen's Printer for Ontario; 2013.
16. Ontario Agency for Health Protection and Promotion (Public Health Ontario). Vector-borne diseases 2012 summary report. Toronto, ON: Queen's Printer for Ontario; 2013.
17. Public Health Division Ministry of Health and Long-Term Care. West Nile virus preparedness and prevention plan 2008. Toronto, ON; 2008.
18. Scott TW, Weaver SC. Eastern equine encephalomyelitis: epidemiology and evolution of mosquito transmission. *Adv Virus Res.* 1989;37:277-328.
19. Smithburn KC, Hughes TP, Burke AW, et al. A neurotropic virus isolated from the blood of a native of Uganda. *Am J Trop Med Hyg.* 1940;20:471-472.
20. Solomon T. Flavivirus encephalitis. *N Engl J Med.* 2004;351(4):370-378.
21. Weaver SC, Barrett ADT. Transmission cycles, host range, evolution and emergence of arboviral disease. *Nat Rev Micro.* 2004;2(10):789-801.